

Lecture 3: Properties of Estimators

Bias · Variance · MSE · Consistency · Sufficiency · Cramér–Rao

We use estimators every day. Are they any good?


We already use estimators (Lecture 1, plug-in principle):

$$\bar{X} \text{ for } \mu, \quad S^2 \text{ for } \sigma^2, \quad \hat{p} = \frac{\text{count}}{n} \text{ for } p$$

But how do we judge an estimator?

Is it close to the truth? How much does it jump around? Can we do better?

$$\frac{(X - \bar{X})^2}{n-1}$$



$$\mu = E[\bar{X}]$$
$$\bar{X} + 1$$


Bias: Is the Estimator Centered on the Truth?

$$\text{Bias}(\hat{\theta}) = \mathbb{E}[\hat{\theta}] - \theta$$


θ

$\mathbb{E}[\hat{\theta}]$

Unbiased



Biased



If $\text{Bias}(\hat{\theta}) = 0$ for all θ , the estimator is **unbiased**.

Worked Example: Is $\bar{X}$ Unbiased for μ ?

Let $X_1, \dots, X_n$ be i.i.d. with $\mathbb{E}[X_i] = \mu$. Is $\hat{\mu} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ unbiased?

Step 1: Compute $\mathbb{E}[\hat{\mu}]$:

$$\mathbb{E}[\bar{X}] = \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i] = \frac{1}{n} \cdot n\mu = \mu$$

Worked Example: Is $\bar{X}$ Unbiased for μ ?

Let $X_1, \dots, X_n$ be i.i.d. with $\mathbb{E}[X_i] = \mu$. Is $\hat{\mu} = \bar{X} = \frac{1}{n} \sum_{i=1}^n X_i$ unbiased?

Step 1: Compute $\mathbb{E}[\hat{\mu}]$:

$$\mathbb{E}[\bar{X}] = \mathbb{E}\left[\frac{1}{n} \sum_{i=1}^n X_i\right] = \frac{1}{n} \sum_{i=1}^n \mathbb{E}[X_i] = \frac{1}{n} \cdot n\mu = \mu$$

Step 2: Check bias:

$$\text{Bias}(\bar{X}) = \mathbb{E}[\bar{X}] - \mu = \mu - \mu = 0$$

✓ Unbiased!

Recipe for any estimator:

- (1) Compute $\mathbb{E}[\hat{\theta}]$ → (2) Subtract the true θ → (3) If the result is 0, it's unbiased.

Worked Example: Why Dividing by n Is Biased

We want to estimate $\sigma^2 = \text{Var}(X_i)$. Natural guess: $\hat{\sigma}_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$.

Trick: rewrite each $(X_i - \bar{X})$ by adding and subtracting the true mean μ :

$$\underbrace{X_i - \bar{X}} = \underbrace{(X_i - \mu)}_{\text{deviation from truth}} - \underbrace{(\bar{X} - \mu)}_{\text{estimation error}}$$

Squaring and summing gives the **key identity**:

$$\sum_{i=1}^n (X_i - \bar{X})^2 = \sum_{i=1}^n (X_i - \mu)^2 - n(\bar{X} - \mu)^2$$

$$\sigma^2 = E(X - E_X)^2$$
$$\frac{1}{n} (X_i - \bar{X})^2$$

Worked Example: Why Dividing by n Is Biased

We want to estimate $\sigma^2 = \text{Var}(X_i)$. Natural guess: $\hat{\sigma}_n^2 = \frac{1}{n} \sum_{i=1}^n (X_i - \bar{X})^2$.

Trick: rewrite each $(X_i - \bar{X})$ by adding and subtracting the true mean μ :

$$X_i - \bar{X} = \underbrace{(X_i - \mu)}_{\text{deviation from truth}} - \underbrace{(\bar{X} - \mu)}_{\text{estimation error}}$$

Squaring and summing gives the **key identity**:

$$\sum_{i=1}^n (X_i - \bar{X})^2 = \sum_{i=1}^n (X_i - \mu)^2 - n(\bar{X} - \mu)^2$$

Take expectations (using $\mathbb{E}[(X_i - \mu)^2] = \sigma^2$ and $\text{Var}(\bar{X}) = \sigma^2/n$):

$$\mathbb{E}\left[\sum (X_i - \mu)^2\right] = n\sigma^2$$

(n terms, each σ^2)

$$\mathbb{E}[n(\bar{X} - \mu)^2] = n \cdot \text{Var}(\bar{X}) = n \cdot \frac{\sigma^2}{n} = \sigma^2$$

(one "lost" degree of freedom)

$$\Rightarrow \mathbb{E}\left[\sum (X_i - \bar{X})^2\right] = n\sigma^2 - \sigma^2 = (n-1)\sigma^2$$

$$n\sigma^2 - \sigma^2 = (n-1)\sigma^2$$

$$(\sqrt{\sigma^2})$$

$$E(\bar{X} - \mu)^2$$

$$\begin{aligned} V, \bar{X} &= \\ &= \text{Var} \frac{1}{n} X_1 + \dots + X_n \\ &= \frac{1}{n^2} (\text{Var } X_1) \cdot n \end{aligned}$$

$$\frac{\sigma^2 \cdot n}{n^2}$$

Bessel's Correction: The Fix

From the previous slide: $\mathbb{E}[\sum (X_i - \bar{X})^2] = (n-1)\sigma^2$, so:

$$\mathbb{E}[\hat{\sigma}_n^2] = \mathbb{E}\left[\frac{1}{n} \sum (X_i - \bar{X})^2\right] = \frac{(n-1)\sigma^2}{n} \neq \sigma^2 \quad \text{Biased!}$$

It **underestimates** by σ^2/n . Why? We used $\bar{X}$ instead of μ , "using up" one degree of freedom.

Bessel's correction: Divide by $n-1$ instead of n :

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2 \quad \mathbb{E}[S^2] = \sigma^2 \quad \checkmark \text{ Unbiased!}$$



Intuition: We estimated μ from the same data, so the residuals $(X_i - \bar{X})$ are "too small" on average. Dividing by $n-1$ corrects for this.

Bias: Summary

$$(X_1, X_2, \dots) \sim \mathcal{N}$$

$$\bar{X} \sim \mathcal{N}$$

Estimator	Bias	Unbiased?
$\bar{X} = \frac{1}{n} \sum X_i$ for μ	0	Yes
$\hat{\sigma}_n^2 = \frac{1}{n} \sum (X_i - \bar{X})^2$ for σ^2	$-\frac{\sigma^2}{n}$	No
$S^2 = \frac{1}{n-1} \sum (X_i - \bar{X})^2$ for σ^2	0	Yes
$\hat{p} = \frac{\sum X_i}{n}$ for p (Bernoulli)	0	Yes



$$E(X_1)$$

Dividing by n instead of $n-1$ **underestimates** the true variance. Bessel's correction ($n-1$) fixes this. Recall Lecture 2!

$$E(\hat{\theta}) = \theta$$




$$X_1, \dots, X_n$$

Unbiasedness Alone Isn't Enough

Consider estimating $\mu = \mathbb{E}[X_i]$ from $X_1, \dots, X_n$.

Surprising fact: $\tilde{\mu} = X_1$ is also **unbiased**!

$$\mathbb{E}[X_1] = \mu \quad \Rightarrow \quad \text{Bias}(X_1) = 0 \quad \checkmark$$



Unbiasedness Alone Isn't Enough

Consider estimating $\mu = \mathbb{E}[X_i]$ from $X_1, \dots, X_n$.

Surprising fact: $\tilde{\mu} = X_1$ is also **unbiased**!

$$\mathbb{E}[X_1] = \mu \Rightarrow \text{Bias}(X_1) = 0 \quad \checkmark$$

But it's a terrible estimator — it ignores $X_2, \dots, X_n$ entirely!

Three statisticians go deer hunting.

The first one shoots **1 meter to the left** of the deer.

The second one shoots **1 meter to the right**.

The third one shouts: "We got it!"

On average they hit the target — **unbiased**! But not very useful...


Lesson: Unbiasedness only says $\mathbb{E}[\hat{\theta}] = \theta$. It says nothing about how much $\hat{\theta}$ **varies**. We need more: **variance** and **MSE**.



$$\tilde{\theta} \rightarrow \mu$$



The Dartboard Analogy



Bullseye = true θ . Blue dots = estimates from repeated samples.

Variance of an Estimator

The **variance** measures how much $\hat{\theta}$ wobbles across samples: $\text{Var}(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \mathbb{E}[\hat{\theta}])^2]$

Why is $\text{Var}(\bar{X}) = \sigma^2/n$ and not σ^2/n^2 ?

Var gives

($\frac{1}{n}$ comes out as $\frac{1}{n^2}$)

$$\text{Var}(\bar{X}) = \text{Var}\left(\frac{1}{n} \sum_{i=1}^n X_i\right) = \frac{1}{n^2} \text{Var}\left(\sum_{i=1}^n X_i\right)$$

$$= \frac{1}{n^2} \sum_{i=1}^n \text{Var}(X_i)$$

(independent $\Rightarrow$ variances **add**)

$$= \frac{1}{n^2} \cdot n\sigma^2 = \boxed{\frac{\sigma^2}{n}}$$

$$\mathbb{E}(\hat{\theta} - \theta)^2$$

(n terms cancel one n)

$$\text{SE}(\bar{X}) = \frac{\sigma}{\sqrt{n}} \quad (\text{standard error} = \sqrt{\text{Var}})$$

~~X~~

Mean Squared Error: The Total Error

Bias tells us about the **aim**, variance about the **spread**. Can we combine them?

Mean Squared Error:

$$\text{MSE}(\hat{\theta}) = \mathbb{E}[(\hat{\theta} - \theta)^2]$$

The average squared distance from the estimate to the truth.

The trick: add and subtract $\mathbb{E}[\hat{\theta}]$ to decompose the error:

$$\hat{\theta} - \theta = \underbrace{(\hat{\theta} - \mathbb{E}[\hat{\theta}])}_{\text{random fluctuation}} + \underbrace{(\mathbb{E}[\hat{\theta}] - \theta)}_{\text{bias (a constant!)}}$$

This splits the total error into two pieces: the **random part** (how much $\hat{\theta}$ moves around its own mean) and the **systematic part** (how far that mean is from the truth).

MSE = Bias² + Variance: The Proof

Now square $\hat{\theta} - \theta = (\hat{\theta} - \mathbb{E}[\hat{\theta}]) + (\mathbb{E}[\hat{\theta}] - \theta)$ and take expectations:

$$\mathbb{E}[(\hat{\theta} - \theta)^2] = \mathbb{E}[(\hat{\theta} - \mathbb{E}[\hat{\theta}])^2] + 2 \underbrace{(\mathbb{E}[\hat{\theta}] - \theta)}_{\text{constant}} \underbrace{\mathbb{E}[\hat{\theta} - \mathbb{E}[\hat{\theta}]]}_{= 0 \text{ (always!)}} + (\mathbb{E}[\hat{\theta}] - \theta)^2$$

The cross term vanishes because $\hat{\theta} - \mathbb{E}[\hat{\theta}]$ has mean zero **by definition**.

$$\text{MSE}(\hat{\theta}) = \text{Var}(\hat{\theta}) + \text{Bias}^2(\hat{\theta})$$



Unbiased means $\text{MSE} = \text{Var}$, but a biased estimator can still win if its variance is low enough.

When Biased Beats Unbiased

Example: Estimating σ^2 from $X_1, \dots, X_n \sim N(\mu, \sigma^2)$.

Estimator	Bias	Variance	MSE
$S^2 = \frac{1}{n-1} \sum (X_i - \bar{X})^2$	0	$\frac{2\sigma^4}{n-1}$	$\frac{2\sigma^4}{n-1}$
$\hat{\sigma}_n^2 = \frac{1}{n} \sum (X_i - \bar{X})^2$	$-\frac{\sigma^2}{n}$	$\frac{2(n-1)\sigma^4}{n^2}$	$\frac{(2n-1)\sigma^4}{n^2}$

Handwritten notes:

- G^4
- $\frac{2}{n-1}$
- $\frac{2n-1}{n^2}$

Compare: $\frac{2n-1}{n^2}$ vs $\frac{2}{n-1} \Rightarrow \hat{\sigma}_n^2$ has **lower MSE** for all $n \geq 2$!


The biased estimator beats the unbiased S^2 because its variance reduction outweighs the small bias.

The Bias-Variance Tradeoff

You can't minimize bias and variance at the same time.

How do we find the **sweet spot**?

The Bias-Variance Tradeoff



Bias-Variance in Machine Learning

This tradeoff is **everywhere** in ML — it's the same principle in different disguises:

Setting	Too simple (high bias)	Too <u>complex</u> (high var)
Polynomial regression	<u>Degree 1 (line)</u>	Degree 20 (wiggly)
KNN	Large k (oversmoothed)	$k = 1$ (memorizes noise)
Decision tree	Shallow tree (underfits)	Deep tree (overfits)
Neural network	Too few neurons	Too many neurons
Regularization	Strong penalty (λ large)	No penalty ($\lambda = 0$)

Key insight: In all these cases, the total error (MSE, test loss) is minimized at an intermediate complexity. This is why we need **cross-validation**, **regularization**, and **held-out test sets** — to find the sweet spot empirically.

bin

X_1

$$E[X_1] + \epsilon = 0$$

Consistency

Does our estimator converge to the truth
as we collect more and more data?

$\bar{X}$ $n \rightarrow \infty$


$X_1, X_2, \dots, X_{100}$



Consistency: Getting It Right Eventually

An estimator $\hat{\theta}_n$ is **consistent** if it converges to the truth as $n \rightarrow \infty$:

$$\hat{\theta}_n \xrightarrow{P} \theta \quad \text{i.e.,} \quad \Pr(|\hat{\theta}_n - \theta| > \varepsilon) \rightarrow 0 \text{ for all } \varepsilon > 0$$



Consistent vs Inconsistent: A Contrast

MSE

$\bar{V}_n$

Consistent: $\hat{\mu} = \bar{X}_n$

Not consistent: $\tilde{\mu} = X_1$

- ▶ $\mathbb{E}[\bar{X}_n] = \mu$ (unbiased)
- ▶ $\text{Var}(\bar{X}_n) = \sigma^2/n \rightarrow 0$
- ▶ Uses **all** n observations
- ▶ More data $\Rightarrow$ more precise

$\sigma^2/n \rightarrow 0$

- ▶ $\mathbb{E}[X_1] = \mu$ (also unbiased!)
- ▶ $\text{Var}(X_1) = \sigma^2$ (constant!)
- ▶ Uses **only** the first observation
- ▶ Ignores all other data forever

Unbiased $\neq$ consistent. X_1 is unbiased but NOT consistent.

Consistent $\neq$ unbiased. $\hat{\sigma}_n^2 = \frac{1}{n} \sum (X_i - \bar{X})^2$ is biased but IS consistent
(because its bias $\rightarrow 0$ and its variance $\rightarrow 0$).

$$\frac{1}{n} \sum (x_i - \bar{x})^2$$

Sufficient Conditions for Consistency

$$P(|X - \mu| \geq \epsilon) \leq \frac{\sigma^2}{\epsilon^2}$$

Chebyshev's inequality gives us a concrete tool:

$$\Pr(|\hat{\theta}_n - \theta| \geq \epsilon) \leq \frac{\mathbb{E}[(\hat{\theta}_n - \theta)^2]}{\epsilon^2} = \frac{\text{MSE}(\hat{\theta}_n)}{\epsilon^2} = \frac{\text{Bias}^2 + \text{Var}}{\epsilon^2}$$

$$\text{MSE}(\hat{\theta}_n) = \mathbb{E}[(\hat{\theta}_n - \theta)^2]$$

$$\text{Bias}(\hat{\theta}_n) \rightarrow 0 \text{ as } n \rightarrow \infty$$

$$\text{Var}(\hat{\theta}_n) \rightarrow 0 \text{ as } n \rightarrow \infty$$


$$\text{MSE} \rightarrow 0 \Rightarrow \Pr(|\hat{\theta}_n - \theta| \geq \epsilon) \rightarrow 0 \Rightarrow \text{consistent!}$$

$$\mathbb{E}[(\hat{\theta}_n - \mu)^2]$$

Example: $\bar{X}_n$ is consistent for μ : Bias = 0, Var = $\sigma^2/n \rightarrow 0$, so $\Pr(|\bar{X}_n - \mu| \geq \epsilon) \leq \sigma^2/(n\epsilon^2) \rightarrow 0$.

This is precisely the **(Weak) Law of Large Numbers**: $\bar{X}_n \xrightarrow{P} \mu$

$$\frac{\sigma^2}{n} \mathbb{E}[(X - \mu)^2] = \mathbb{E}[(\hat{\theta}_n - \theta)^2]$$



$\#T$

$T \rightarrow \frac{4}{5}$

Sufficiency

We have n data points. Do we really need **all** of them?

Can we **compress** without losing information?

$$T(x_1, x_2, \dots, x_n) = \# \{x_i = T\} \quad P(T) = \frac{4}{5}$$

Sufficiency: Can We Compress the Data?

Example: $X_1, \dots, X_n \sim \text{Bern}(p)$. To estimate p :

- ▶ We only need $T = \sum X_i$ (total number of successes)
- ▶ The specific order (HHTHT vs THHHTH) tells us nothing more about p

$$0 \ 1 \ 1 \ 0$$
$$T(x_1; \dots; x_n) = \sum_{i=1}^n x_i$$


Definition: A statistic $T(\mathbf{X})$ is **sufficient** for θ if the conditional distribution of $\mathbf{X} \mid T(\mathbf{X})$ does not depend on θ .

Intuition: Once you know T the remaining randomness in the data is just noise — it carries **no information** about θ . T is a “lossless summary.”

Sufficiency as Data Compression

$$T(X) = G$$

$$y(t) = \frac{t(x)}{n}$$



Example

Bernoulli

$$0, 1, 1, 0, 1, 1, 1, 0, 1, 0 \longrightarrow T = \sum X_i = 6 \longrightarrow \hat{p} = 6/10 = 0.6$$

$\uparrow$

The order $(0, 1, 1, 0, 1, \dots)$ doesn't matter for estimating p — only the **total count** matters.

How to Check: Fisher–Neyman Factorization

Theorem: $T(\mathbf{X})$ is sufficient for θ if and only if:

$$f(\mathbf{x} | \theta) = g(T(\mathbf{x}), \theta) \cdot h(\mathbf{x})$$

where g depends on the data **only through** T and h does not depend on θ .

Bernoulli worked example: $X_1, \dots, X_n \sim \text{Bern}(p)$, let $T = \sum X_i$.

$$f(\mathbf{x} | p) = \prod_{i=1}^n p^{x_i} (1-p)^{1-x_i} = \underbrace{p^{\sum x_i} (1-p)^{n-\sum x_i}}_{g(T, p)} \cdot \underbrace{1}_{h(\mathbf{x})}$$

Model

Sufficient statistic

Intuition

Bern(p)

$T = \sum X_i$

1 number for 1 parameter

$N(\mu, \sigma_0^2)$ (σ_0^2 known)

$T = \bar{X}$

1 number for 1 parameter

$N(\mu, \sigma^2)$ (both unknown)

$T = (\bar{X}, S^2)$

2 numbers for 2 parameters

$$f(\mathbf{x} | \theta) =$$

$$= g(T(\mathbf{x}), \theta) \cdot h(\mathbf{x})$$

$$f(x_i | p) = \begin{cases} p & \text{if } x_i = 1 \\ 1-p & \text{if } x_i = 0 \end{cases}$$

$$p^{x_i} (1-p)^{1-x_i}$$

$$T(\mathbf{x}) = \sum x_i$$

$$T(\mathbf{x}) = \begin{bmatrix} \sum x_i \\ \sum (1-x_i) \end{bmatrix}$$

Minimal Sufficiency

$$T(X_1, X_2, \dots, X_n) = (\bar{x}, s^2)$$

The full data $\mathbf{X}$ is always trivially sufficient. But can we compress **further**?

$$\sum x_i$$

$$p(y_1, y_2, y_3) = y_1$$

$$T(x) = \sum x_i$$

A sufficient statistic is minimal if it is a function of every other sufficient statistic.

It achieves the **maximum compression** without losing information about θ .

$$\varphi(x_1, x_2, \dots, x_n) = T(x)$$

Example: For $X_1, \dots, X_n \sim N(\mu, \sigma_0^2)$ with σ_0^2 known:

$\mathbf{X} = (X_1, \dots, X_n)$
sufficient, no compression

compress

$(\bar{X}, S^2)$
sufficient, partial

compress more

$\bar{X}$ alone
sufficient and minimal

Since only μ is unknown, S^2 carries no extra information — $\bar{X}$ alone is enough.

The Rao-Blackwell Theorem

Why does sufficiency matter for estimation? Because it lets us **improve** any estimator:

Rao-Blackwell Theorem: Given *any* unbiased estimator $\tilde{\theta}$ and a sufficient statistic T , define $\hat{\theta} = \mathbb{E}[\tilde{\theta} \mid T]$. Then:

- (1) $\hat{\theta}$ is still unbiased: $\mathbb{E}[\hat{\theta}] = \mathbb{E}[\tilde{\theta}] = \theta$
- (2) $\hat{\theta}$ has **lower or equal variance**: $\text{Var}(\hat{\theta}) \leq \text{Var}(\tilde{\theta})$

Conditioning on a sufficient statistic **never hurts, often helps**.

$\tilde{\theta}$
+

$$\hat{\theta} = \mathbb{E}[\tilde{\theta} \mid T]$$

6-20
17.39210w

The Rao-Blackwell Theorem

Why does sufficiency matter for estimation? Because it lets us **improve** any estimator:

Rao-Blackwell Theorem: Given *any* unbiased estimator $\tilde{\theta}$ and a sufficient statistic T , define $\hat{\theta} = \mathbb{E}[\tilde{\theta} \mid T]$. Then:

- (1) $\hat{\theta}$ is still **unbiased**: $\mathbb{E}[\hat{\theta}] = \mathbb{E}[\tilde{\theta}] = \theta$
- (2) $\hat{\theta}$ has **lower or equal variance**: $\text{Var}(\hat{\theta}) \leq \text{Var}(\tilde{\theta})$

Conditioning on a sufficient statistic **never hurts, often helps**.

Worked example: $X_1, \dots, X_n \sim \text{Bern}(p)$, sufficient stat $T = \sum X_i$.

$$\underbrace{\tilde{p} = X_1}_{\text{naive: unbiased, Var}=p(1-p)} \xrightarrow{\mathbb{E}[\cdot | T]} \underbrace{\hat{p} = \mathbb{E}[X_1 | T] = T/n = \bar{X}}_{\text{improved: unbiased, Var}=p(1-p)/n} \quad \times n \text{ better!}$$

What Does $\mathbb{E}[\tilde{\theta} \mid T]$ Actually Mean?

Concrete example: $X_1, X_2, X_3 \sim \text{Bern}(p)$, $T = X_1 + X_2 + X_3$, $\tilde{p} = X_1$.

Suppose someone tells you $T = 2$ (two successes). Which data vectors give $T = 2$?

(X_1, X_2, X_3)	T	$\tilde{p} = X_1$	Equally likely?
$(1, 1, 0)$	2	1	Yes
$(1, 0, 1)$	2	1	Yes
$(0, 1, 1)$	2	0	Yes

$$T(x) = 2$$

~~1 1 0~~
~~0 0 1~~
~~0 0 0~~

$$\tilde{\theta} \mid T$$

What Does $\mathbb{E}[\tilde{\theta} \mid T]$ Actually Mean?

Concrete example: $X_1, X_2, X_3 \sim \text{Bern}(p)$, $T = X_1 + X_2 + X_3$, $\tilde{p} = X_1$.

Suppose someone tells you $T = 2$ (two successes). Which data vectors give $T = 2$?

(X_1, X_2, X_3)	T	$\tilde{p} = X_1$	Equally likely?
$(1, 1, 0)$	2	1	Yes
$(1, 0, 1)$	2	1	Yes
$(0, 1, 1)$	2	0	Yes

$$\mathbb{E}[X_1 \mid T = 2] = \frac{1 + 1 + 0}{3} = \frac{2}{3} = \frac{T}{n} \quad \checkmark$$

“Condition on T ” means: average $\tilde{\theta}$ over all data configurations that produce the same value of T . The noise (which specific X_i ’s are 1 vs 0) gets averaged away. Only the useful part (T) survives.

Why Does Rao–Blackwell Work?

The key is the law of total variance (a.k.a. Eve's law):

$$\text{Var}(\tilde{\theta}) = \underbrace{\mathbb{E}[\text{Var}(\tilde{\theta} | T)]}_{\text{"useless" noise} \geq 0} + \text{Var}\left(\underbrace{\mathbb{E}[\tilde{\theta} | T]}_{\hat{\theta}}\right)$$

$$\text{Var}(\hat{\theta} | T) + \mathbb{E}(\tilde{\theta} | T)$$


Why Does Rao–Blackwell Work?

The key is the **law of total variance** (a.k.a. Eve's law):

$$\text{Var}(\tilde{\theta}) = \underbrace{\mathbb{E}[\text{Var}(\tilde{\theta} \mid T)]}_{\text{"useless" noise} \geq 0} + \text{Var}\left(\underbrace{\mathbb{E}[\tilde{\theta} \mid T]}_{\hat{\theta}}\right)$$

Since the first term ≥ 0 , we immediately get:

$$\text{Var}(\tilde{\theta}) \geq \text{Var}(\hat{\theta})$$



Intuition: T captures all the useful information about θ . Conditioning on T removes the "useless" randomness (the part that doesn't tell us about θ). What's left is a cleaner estimator.

Finding Minimal Sufficient Statistics

Theorem (Likelihood Ratio Criterion): $T(\mathbf{X})$ is minimal sufficient iff for all $\mathbf{x}, \mathbf{y}$:

$$\underline{T(\mathbf{x})} = T(\mathbf{y}) \iff \frac{f(\mathbf{x} | \theta)}{f(\mathbf{y} | \theta)} \text{ does not depend on } \theta$$

Bernoulli example: $X_1, \dots, X_n \sim \text{Bern}(p)$.

$$\frac{f(\mathbf{x} | p)}{f(\mathbf{y} | p)} = \frac{p^{\sum x_i} (1-p)^{n-\sum x_i}}{p^{\sum y_i} (1-p)^{n-\sum y_i}} = \left(\frac{p}{1-p} \right)^{\sum x_i - \sum y_i}$$

Free of $p \iff \sum x_i = \sum y_i$. So $T = \sum X_i$ is **minimal sufficient** for p . ✓

Recipe: Write the likelihood ratio $f(\mathbf{x} | \theta)/f(\mathbf{y} | \theta)$.

Find which function of the data must match for the ratio to lose its θ -dependence.

That function is the minimal sufficient statistic.

The Exponential Family: A Unifying Framework

All our examples — Bernoulli, Normal, Poisson, Exponential — share one structure:

$$f(x | \theta) = h(x) \exp(\eta(\theta) T(x) - A(\theta))$$

Distribution	Natural param $\eta(\theta)$	$T(x)$	Suff. stat (n obs)
Bern(p)	$\log \frac{p}{1-p}$	x	$\sum X_i$
$N(\mu, \sigma_0^2)$ (σ_0^2 known)	μ / σ_0^2	x	$\sum X_i$
Pois(λ)	$\log \lambda$	x	$\sum X_i$
Exp(λ)	$-\lambda$	x	$\sum X_i$

Pattern: For single-parameter families, $T(x) = x$. The sufficient statistic for n observations is always $\sum T(X_i)$ — straight from the factorization theorem!

Why Exponential Families Are Special

Nearly every nice property we've discussed is **automatic** in exponential families:

Sufficiency: $T(\mathbf{X}) = \sum T(X_i)$ is sufficient **and minimal**

Completeness: the natural sufficient statistic is **complete** (see below)

Regularity: all conditions for the Cramér–Rao bound (coming soon) are satisfied

Optimal estimators exist: we'll see this when we reach the CR bound

Completeness means: if $\mathbb{E}_\theta[g(T)] = 0$ for all θ , then $g(T) = 0$ a.s. $\rightarrow$ **no non-trivial unbiased estimator of zero** based on T .


Lehmann–Scheffé: An unbiased estimator based on a **complete** sufficient statistic is the **unique best** unbiased estimator (UMVUE). For exp. families, $\sum T(X_i)$ is always complete $\Rightarrow$ UMVUE exists!

Can We Do Better? The Fundamental Question

We know $\text{Var}(\bar{X}) = \sigma^2/n$ for estimating the mean.

Can **any** unbiased estimator have **lower** variance?

Or is $\bar{X}$ already the best we can do?



To answer this, we need to measure **how much information** one observation carries about θ .

Roadmap:

Why log? → **Score function** (sensitivity of the model to θ) → **Fisher information**
→ **Cramér–Rao bound** (the variance floor)

Why the Logarithm? From Products to Sums

The likelihood for i.i.d. data is a **product**:

$$L(\theta) = \prod_{i=1}^n f(X_i | \theta)$$

Taking the log turns this into a **sum**:

$$\ell(\theta) = \log L(\theta) = \sum_{i=1}^n \log f(X_i | \theta)$$

Products are painful:

- ▶ Multiplying tiny numbers $\rightarrow$ underflow
- ▶ Product rule for derivatives is messy
- ▶ Hard to work with analytically

Sums are friendly:

- ▶ Numerically stable
- ▶ Derivative of a sum = sum of derivatives
- ▶ LLN, CLT apply directly

Key fact: log is monotonically increasing, so $\arg \max_{\theta} L(\theta) = \arg \max_{\theta} \ell(\theta)$. Same maximizer!

The Score Function: How Sensitive Is the Model?

Given a model $f(x | \theta)$, the **score** measures how the log-probability changes with θ :

$$s(\theta) = \frac{\partial}{\partial \theta} \log f(X | \theta)$$

Concrete example: $X \sim \text{Bernoulli}(p)$.

$$\log f(x | p) = x \log p + (1-x) \log(1-p)$$

$$s(p) = \frac{\partial}{\partial p} [x \log p + (1-x) \log(1-p)] = \frac{x}{p} - \frac{1-x}{1-p} = \frac{x-p}{p(1-p)}$$

- ▶ If we observe $x = 1$ and p is small, the score is **large positive** $\rightarrow$ “ p should be higher”
- ▶ If we observe $x = 0$ and p is large, the score is **large negative** $\rightarrow$ “ p should be lower”
- ▶ On average: $\mathbb{E}[s(p)] = 0$ — the score points in the right direction but **averages out**

Fisher Information: How Informative Is One Observation?


The score averages to zero, but it **varies**. More variation = more information:

$$I(\theta) = \text{Var}[s(\theta)] = \mathbb{E} \left[\left(\frac{\partial}{\partial \theta} \log f(X | \theta) \right)^2 \right]$$

Bernoulli derivation: We found $s(p) = \frac{X-p}{p(1-p)}$.

Since $\mathbb{E}[s] = 0$:

$$\begin{aligned} I(p) &= \mathbb{E}[s^2] = \mathbb{E} \left[\frac{(X-p)^2}{p^2(1-p)^2} \right] \\ &= \frac{\text{Var}(X)}{p^2(1-p)^2} = \frac{p(1-p)}{p^2(1-p)^2} = \boxed{\frac{1}{p(1-p)}} \end{aligned}$$



p near 0 or 1: very informative. $p = 0.5$: max noise, min info.

Fisher Information: Two Equivalent Forms

Under regularity conditions, there is an equivalent formula that's often easier to compute:

$$I(\theta) = \mathbb{E}[s(\theta)^2] = -\mathbb{E}\left[\frac{\partial^2}{\partial \theta^2} \log f(X | \theta)\right]$$

Why are these the same? Start from $\mathbb{E}[s(\theta)] = 0$ and differentiate both sides w.r.t. θ :



$$0 = \frac{\partial}{\partial \theta} \mathbb{E}[s] = \mathbb{E}\left[\frac{\partial s}{\partial \theta}\right] + \mathbb{E}[s \cdot s] = \mathbb{E}[\ell''] + \mathbb{E}[s^2]$$

So: $\mathbb{E}[s^2] = -\mathbb{E}[\ell'']$. ✓

Verify for Bernoulli: $\ell(p) = x \log p + (1-x) \log(1-p)$

$$\ell''(p) = -\frac{x}{p^2} - \frac{1-x}{(1-p)^2} \Rightarrow -\mathbb{E}[\ell''] = \frac{p}{p^2} + \frac{1-p}{(1-p)^2} = \frac{1}{p} + \frac{1}{1-p} = \frac{1}{p(1-p)} \quad \checkmark$$

Intuition: Sharp vs Flat Log-Likelihood



$I(\theta)$ measures the **curvature** of the log-likelihood at the true θ .

Sharp curve $\Rightarrow$ high $I(\theta)$ $\Rightarrow$ data is very informative $\Rightarrow$ estimator is precise.

This connects the two forms: $I(\theta) = -\mathbb{E}[\ell'']$ is literally the expected curvature.

Cramér–Rao Lower Bound

Now we can answer the fundamental question. For any **unbiased** estimator $\hat{\theta}$ based on n i.i.d. observations:

$$\text{Var}(\hat{\theta}) \geq \frac{1}{n \cdot I(\theta)}$$

Intuition: Why $\frac{1}{n \cdot I(\theta)}$?

- ▶ **More observations (n large)** $\Rightarrow$ bound gets smaller $\Rightarrow$ can estimate more precisely
- ▶ **More informative data ($I(\theta)$ large)** $\Rightarrow$ bound gets smaller $\Rightarrow$ each observation tells us more
- ▶ The bound is **tight** for many models — it's the actual achievable precision

Verify for Bernoulli:

$$I(p) = \frac{1}{p(1-p)} \quad \Rightarrow \quad \text{CR bound: } \text{Var}(\hat{p}) \geq \frac{1}{n \cdot \frac{1}{p(1-p)}} = \frac{p(1-p)}{n}$$

Actual variance of $\hat{p} = \bar{X}$: $\text{Var}(\hat{p}) = \frac{p(1-p)}{n}$ ✓ Hits the bound exactly!

Cramér–Rao: Efficiency and Practical Use

What it says:

There is a **floor** on how precise any unbiased estimator can be

Efficient estimator:

Achieves the bound – the **best possible**

Practical use:

Tells you whether to keep searching for a better estimator

Model	Estimator	$\text{Var}(\hat{\theta})$	CR bound	Efficient?
$\text{Bern}(p)$	$\hat{p} = \bar{X}$	$\frac{p(1-p)}{n}$	$\frac{p(1-p)}{n}$	Yes
$N(\mu, \sigma_0^2)$	$\hat{\mu} = \bar{X}$	$\frac{\sigma_0^2}{n}$	$\frac{\sigma_0^2}{n}$	Yes
$\text{Exp}(\lambda)$	$\hat{\lambda} = 1/\bar{X}$	$\frac{\lambda^2}{n}$	$\frac{\lambda^2}{n}$	Yes


Regularity Conditions: When Does CR Apply?

The Cramér–Rao bound requires **regularity conditions**:

1. **Support** of $f(x | \theta)$ doesn't depend on θ
2. θ in the **interior** of the parameter space
3. Can **differentiate under the integral** sign (swap $\frac{\partial}{\partial \theta}$ and $\int$)
4. $0 < I(\theta) < \infty$ (finite, positive info)

Counterexample: Uniform(0, θ)

- ▶ Support $[0, \theta]$ depends on θ ! (violates #1)
- ▶ Suff. stat: $X_{(n)} = \max_i X_i$
- ▶ $\text{Var}(X_{(n)}) \sim 1/n^2$ — **faster** than CR!
(CR would give $1/n$, but $1/n^2$ is possible here)




Good news: All exponential family distributions automatically satisfy the regularity conditions. The CR bound always applies to them.

Admissibility

Definition: $\hat{\theta}_1$ is **inadmissible** if $\exists \hat{\theta}_2$ that **dominates** it:

$$\text{MSE}(\hat{\theta}_2, \theta) \leq \text{MSE}(\hat{\theta}_1, \theta) \quad \forall \theta, \quad \text{with strict inequality for some } \theta$$

An estimator is **admissible** if no other estimator dominates it.



$\hat{\theta}_1$ is **inadmissible** — $\hat{\theta}_2$ is at least as good everywhere, and strictly better somewhere.

Stein's Paradox (1956)


Surprising fact:

When estimating $\mu = (\mu_1, \dots, \mu_d)$ from $X_i \sim N(\mu_i, 1)$, the sample mean $\hat{\mu}_i = X_i$ is **inadmissible** when $d \geq 3$!

The **James–Stein estimator** dominates it:

$$\hat{\mu}_i^{JS} = \left(1 - \frac{d-2}{\|\mathbf{X}\|^2}\right) X_i$$


- ▶ **Shrinks** each X_i toward 0
- ▶ Works even if μ_i 's are unrelated!
- ▶ A little bias buys a lot of variance reduction



Paradox: estimating the average temperature in Yerevan *improves* if you jointly estimate it with the price of tea in China and the height of the Eiffel Tower.

Why Does Stein's Paradox Work?

The **MSE comparison** tells the whole story:



Why $d \geq 3$? The shrinkage factor $\frac{d-2}{\|\mathbf{X}\|^2}$ needs to be estimated from data.

- ▶ In $d = 1$ or 2 : not enough “room” — estimation error of the shrinkage factor wipes out the gain
- ▶ In $d \geq 3$: $\|\mathbf{X}\|^2$ concentrates well enough $\rightarrow$ shrinkage factor is accurate $\rightarrow$ net win


Connection to ML: James–Stein shrinkage is an early form of **regularization**.

Ridge regression (L^2 penalty) does the same thing: shrink coefficients toward zero. The bias-variance tradeoff in action: a little bias buys a lot of variance reduction.

Minimax Estimators

A **minimax** estimator minimizes the **worst-case** risk:

$$\hat{\theta}_{\text{minimax}} = \arg \min_{\hat{\theta}} \max_{\theta} \text{MSE}(\hat{\theta}, \theta)$$



Minimax = **conservative**: protects against the worst θ . Minimax hedges.

Three Philosophies of Estimation

Plug-in (unbiased)

Use sample statistic directly
 $(\bar{X}, S^2, \hat{p})$

Admissible in $d = 1$

Inadmissible in $d \geq 3$

Shrinkage

Pull estimates toward a
central value (e.g. 0)

Biased but lower MSE
(James–Stein)

Minimax

Minimize worst-case risk

Conservative guarantee
No single θ can
hurt you badly

Takeaway: In high dimensions ($d \geq 3$), shrinkage estimators are provably better than using each sample statistic on its own. We'll see more of this in later lectures.

What We Haven't Covered (Yet)

This lecture focused on **point estimation** — producing a single “best guess” for θ . But there's much more to statistical inference:

Confidence intervals: How uncertain is our estimate? (Lectures 5–6)

Hypothesis testing: Is the effect real or just noise? (Lectures 7–8)

Bayesian estimation: Incorporating prior beliefs (Lecture 5)

Bootstrap: Resampling to estimate uncertainty without formulas (Lecture 6)

Asymptotic theory: What happens as $n \rightarrow \infty$ in general? (Lecture 5)

Nonparametric estimation: What if we don't assume a distribution at all?

Today's tools (bias, MSE, CR bound, sufficiency) will be the **foundation** for all of these.

Summary: How to Judge an Estimator

Bias: $\mathbb{E}[\hat{\theta}] - \theta$. Does it aim at the right place?

Variance: $\text{Var}(\hat{\theta})$. How much does it jump around?

MSE = $\text{Bias}^2 + \text{Var}$. Total error. Biased can beat unbiased!

Consistency: $\hat{\theta}_n \xrightarrow{P} \theta$. Converges to truth with enough data.

Sufficiency: $T(\mathbf{X})$ captures everything about θ . Compress without loss.

Cramér–Rao: $\text{Var} \geq 1/(n \cdot I(\theta))$. The efficiency floor.

Admissibility: No other estimator dominates it everywhere.

Minimax: Best worst-case guarantee. Shrinkage often wins.

Homework

1. Show that $\bar{X}$ is unbiased for μ and compute its MSE.
2. Show that $\hat{\sigma}_n^2 = \frac{1}{n} \sum (X_i - \bar{X})^2$ is biased for σ^2 . Find the bias.
3. Compute the Fisher information $I(\theta)$ for $\text{Poisson}(\lambda)$.
Use it to find the Cramér–Rao lower bound for estimating λ .
Is $\hat{\lambda} = \bar{X}$ efficient?
4. Suppose you shrink $\bar{X}$ toward 0: $\hat{\mu}_c = c\bar{X}$ for $0 < c < 1$.
Find the bias, variance, and MSE as functions of c .
For what value of c is MSE minimized? Is the optimal estimator biased?
5. Use the factorization theorem to show that $T = \sum X_i$ is a sufficient statistic for λ when $X_1, \dots, X_n \sim \text{Poisson}(\lambda)$.

Questions?